

RESTRICTED USE PESTICIDE

DUE TO HAZARD TO NONTARGET ORGANISMS

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certificate.

FIFRA Section 24(c) Special Local Need (SLN) Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

rozol[®] VOLE BAIT

**FOR AERIAL BROADCAST APPLICATION TO CONTROL MEADOW OR
MOUNTAIN VOLES IN POME FRUIT
(APPLE, PEAR) AND STONE FRUIT (APRICOT, CHERRY, NECTARINE,
PEACH, PLUM, PRUNE) TREE ORCHARDS AFTER FALL HARVEST**

EPA Reg. No. 7173-242 EPA SLN No. WA-070019

This SLN label for Rozol Vole Bait expires and may be amended or cancelled before it expires. Do not distribute or use this SLN label after December 31, 2029, or after this SLN label is amended or cancelled, whichever event happens first.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions and precautions on this SLN label and the EPA registered label.

DIRECTIONS FOR USE:

POME AND STONE FRUIT TREE ORCHARDS FOR MEADOW OR MOUNTAIN VOLE CONTROL:

APPLICATION DIRECTIONS: Before application, examine orchard floor to locate trails and runway systems to be treated. Apply bait when three days of rain-free and snow-free weather are expected. Apply bait after harvest (including all harvestable fruit drops) and before new growth in the spring.

Aerial broadcast baiting: Uniformly apply 10 pounds of bait per acre. If populations are high, make a second application 1 to 2 months after the first. Do not apply more than 20 pounds of bait per acre per season. To avoid exposing nontarget organisms, clean up any bait in piles for reuse or disposal according to the federal label.

RESTRICTIONS / PRECAUTIONS:

- Do not apply more than 10 pounds of bait per acre per application.
- Do not apply more than 20 pounds of bait per acre per year.
- Do not apply bait during the growing season.
- Do not apply bait over bare ground or to crops not specified above.
- Do not expose humans (including children), pets, livestock or other nontarget animals to bait during application, handling, storage or disposal. Do not allow livestock to graze or clean up fruit drops in treated areas. Do not use hay cut after application for livestock feed or bedding.
- Do not apply by air within 50 feet of any body of surface water or where raptors are actively feeding on voles.
- This product is extremely toxic to fish and wildlife. Dogs and predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten the bait. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.
- Rozol Vole Bait should not be used under this SLN label where impact on threatened or endangered species is likely. Consult the federal label for additional restrictions and precautions to protect endangered species.

WSDA Container Disposal Guidance:

Completely empty the contents of the container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

Section 24(c) SLN registrant:

Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209

Rev. 06/03/2025